

ENTIRE SOLUTIONS OF SUBLINEAR ELLIPTIC EQUATIONS IN ANISOTROPIC MEDIA

Teodora-Liliana Dinu

Department of Mathematics, “Frații Buzești” College, 200352 Craiova, Romania

E-mail: tldinu@gmail.com

ABSTRACT. We study the nonlinear elliptic problem $-\Delta u = \rho(x)f(u)$ in $\mathbb{R}^N$ ($N \geq 3$), $\lim_{|x| \rightarrow \infty} u(x) = \ell$, where $\ell \geq 0$ is a real number, $\rho(x)$ is a nonnegative potential belonging to a certain Kato class, and $f(u)$ has a sublinear growth. We distinguish the cases $\ell > 0$ and $\ell = 0$ and we prove existence and uniqueness results if the potential $\rho(x)$ decays fast enough at infinity. Our arguments rely on comparison techniques and on a theorem of Brezis and Oswald for sublinear elliptic equations.

2000 Mathematics Subject Classification: 35B40, 35B50, 35J60, 58J05.

Key words: sublinear elliptic equation, entire solution, maximum principle, anisotropic potential, Kato class.

1 Introduction and main results

In their celebrated paper [3], Brezis and Kamin have been concerned with various questions related to the existence of bounded solutions of the sublinear elliptic equation without condition at infinity

$$-\Delta u = \rho(x)u^\alpha \quad \text{in } \mathbb{R}^N, \quad N \geq 3, \quad (1)$$

where $0 < \alpha < 1$, $\rho \in L_{\text{loc}}^\infty(\mathbb{R}^N)$, $\rho \geq 0$ and $\rho \not\equiv 0$. We summarize in what follows the main results obtained in [3]. Brezis and Kamin proved that the *nonlinear* problem (1) has a bounded solution $u > 0$ if and only if the *linear* problem

$$-\Delta u = \rho(x) \quad \text{in } \mathbb{R}^N$$

has a bounded solution. In this case, Problem (1) has a minimal positive solution and this solution satisfies $\liminf_{|x| \rightarrow \infty} u(x) = 0$. Moreover, the minimal solution is the unique positive solution of (1) which tends to zero at infinity. Brezis and Kamin also showed that if the potential $\rho(x)$ decays fast enough at infinity then Problem (1) has a solution and, moreover, such a solution does not exist if $\rho(x)$ has a slow decay at infinity. For instance, if $\rho(x) = (1 + |x|^p)^{-1}$, then (1) has a bounded solution if and only if $p > 2$. More generally, Brezis and Kamin have proved that Problem (1) has a bounded solution if and only if $\rho(x)$ is potentially bounded, that is, the mapping $x \mapsto \int_{\mathbb{R}^N} \rho(y)|x-y|^{2-N} dy \in L^\infty(\mathbb{R}^N)$. We refer to [4, 12] for various results on bounded domains for sublinear elliptic equations with zero Dirichlet boundary condition. Problem (1) in the whole space has been considered in [5, 6, 8, 9, 11, 13, 14] under various assumptions on ρ . Sublinear problems (either stationary or evolution ones) appear in the study of population dynamics, of reaction-diffusion processes, of filtration in porous media with absorption, as well as in the study of the scalar curvature of warped products of semi-Riemannian manifolds (see, e.g., [16]).

Our purpose in this paper is to study the problem

$$\begin{cases} -\Delta u = \rho(x)f(u) & \text{in } \mathbb{R}^N \\ u > \ell & \text{in } \mathbb{R}^N \\ u(x) \rightarrow \ell & \text{as } |x| \rightarrow \infty, \end{cases} \quad (2)$$

where $N \geq 3$ and $\ell \geq 0$ is a real number.

Throughout the paper we assume that the variable potential $\rho(x)$ satisfies $\rho \in L_{\text{loc}}^\infty(\mathbb{R}^N)$, $\rho \geq 0$ and $\rho \not\equiv 0$.

In our first result we suppose that the growth at infinity of the anisotropic potential $\rho(x)$ is given by

($\rho 1$) $\int_0^\infty r \Phi(r) dr < \infty$, where $\Phi(r) := \max_{|x|=r} \rho(x)$.

Assumption ($\rho 1$) has been first introduced in Naito [14].

The nonlinearity $f : (0, \infty) \rightarrow (0, \infty)$ satisfies $f \in C_{\text{loc}}^{0,\alpha}(0, \infty)$ ($0 < \alpha < 1$) and has a sublinear growth, in the sense that

(f1) the mapping $u \mapsto f(u)/u$ is decreasing on $(0, \infty)$ and $\lim_{u \rightarrow \infty} f(u)/u = 0$.

We point out that condition (f1) does not require that f is smooth at the origin. The standard example of such a nonlinearity is $f(u) = u^p$, where $-\infty < p < 1$. We also observe that we study an equation of the same type as in Brezis and Kamin [3]. The main difference is that we require a certain asymptotic behaviour at infinity of the solution.

Under the above hypotheses ($\rho 1$) and (f1), our first result concerns the case $\ell > 0$. We have

Theorem 1. *Assume that $\ell > 0$. Then Problem (2) has a unique classical solution.*

Next, consider the case $\ell = 0$. Instead of ($\rho 1$) we impose the stronger condition

($\rho 2$) $\int_0^\infty r^{N-1} \Phi(r) dr < \infty$.

We remark that in Edelson [7] it is used the stronger assumption $\int_0^\infty r^{N-1+\lambda(N-2)} \Phi(r) dr < \infty$, for some $\lambda \in (0, 1)$.

Additionally, we suppose that

(f2) f is increasing in $(0, \infty)$ and $\lim_{u \searrow 0} f(u)/u = +\infty$.

A nonlinearity satisfying both (f1) and (f2) is $f(u) = u^p$, where $0 < p < 1$.

Our result in the case $\ell = 0$ is the following.

Theorem 2. *Assume that $\ell = 0$ and assumptions ($\rho 2$), (f1) and (f2) are fulfilled. Then Problem (2) has a unique classical solution.*

We point out that assumptions ($\rho 1$) and ($\rho 2$) are related to a celebrated class introduced by Kato, with wide and deep applications in Potential Theory and Brownian Motion. We recall (see [1]) that a real-valued measurable function ψ on $\mathbb{R}^N$ belongs to the Kato class $\mathcal{K}$ provided that

$$\lim_{\alpha \rightarrow 0} \sup_{x \in \mathbb{R}^N} \int_{|x-y| \leq \alpha} E(y) |\psi(y)| dy = 0,$$

where E denotes the fundamental solution of the Laplace equation. According to this definition and our assumption ($\rho 1$) (resp., ($\rho 2$)), it follows that $\psi = \psi(|x|) \in \mathcal{K}$, where $\psi(|x|) := |x|^{N-3} \Phi(|x|)$ (resp., $\psi(|x|) := |x|^{-1} \Phi(|x|)$), for all $x \neq 0$.

2 Proof of Theorem 1

In order to prove the existence of a solution to Problem (2), we use a result established by Brezis and Oswald (see [4, Theorem 1]) for bounded domains. Consider the problem

$$\begin{cases} -\Delta u = g(x, u) & \text{in } \Omega \\ u \geq 0, \quad u \not\equiv 0 & \text{in } \Omega \\ u = 0 & \text{on } \partial\Omega, \end{cases} \quad (3)$$

where $\Omega \subset \mathbb{R}^N$ is a bounded domain with smooth boundary and $g(x, u) : \Omega \times [0, \infty) \rightarrow \mathbb{R}$. Assume that

$$\begin{cases} \text{for a.e. } x \in \Omega \text{ the function } u \mapsto g(x, u) \text{ is continuous on } [0, \infty) \\ \text{and the mapping } u \mapsto g(x, u)/u \text{ is decreasing on } (0, \infty); \end{cases} \quad (4)$$

$$\text{for each } u \geq 0 \text{ the function } x \mapsto g(x, u) \text{ belongs to } L^\infty(\Omega); \quad (5)$$

$$\exists C > 0 \text{ such that } g(x, u) \leq C(u + 1) \text{ a.e. } x \in \Omega, \quad \forall u \geq 0. \quad (6)$$

Set

$$a_0(x) = \lim_{u \searrow 0} g(x, u)/u \quad \text{and} \quad a_\infty(x) = \lim_{u \rightarrow \infty} g(x, u)/u,$$

so that $-\infty < a_0(x) \leq +\infty$ and $-\infty \leq a_\infty(x) < +\infty$.

Under these hypotheses, Brezis and Oswald proved in [4] that Problem (3) has at most one solution. Moreover, a solution of (3) exists if and only if

$$\lambda_1(-\Delta - a_0(x)) < 0 \quad (7)$$

and

$$\lambda_1(-\Delta - a_\infty(x)) > 0, \quad (8)$$

where $\lambda_1(-\Delta - a(x))$ denotes the first eigenvalue of the operator $-\Delta - a(x)$ with zero Dirichlet condition. The precise meaning of $\lambda_1(-\Delta - a(x))$ is

$$\lambda_1(-\Delta - a(x)) = \inf_{\varphi \in H_0^1(\Omega), \|\varphi\|_{L^2(\Omega)}=1} \left(\int |\nabla \varphi|^2 - \int_{[\varphi \neq 0]} a \varphi^2 \right).$$

Note that $\int_{[\varphi \neq 0]} a \varphi^2$ makes sense if $a(x)$ is any measurable function such that either $a(x) \leq C$ or $a(x) \geq -C$ a.e. on Ω .

For any positive integer k we consider the problem

$$\begin{cases} -\Delta u_k = \rho(x)f(u_k), & \text{if } |x| < k \\ u_k > \ell, & \text{if } |x| < k \\ u_k(x) = \ell, & \text{if } |x| = k. \end{cases} \quad (9)$$

Equivalently, the above boundary value problem can be rewritten

$$\begin{cases} -\Delta v_k = \rho(x)f(v_k + \ell), & \text{if } |x| < k \\ v_k(x) = 0, & \text{if } |x| = k. \end{cases} \quad (10)$$

In order to obtain a solution of the problem (10), it is enough to check the hypotheses of the Brezis-Oswald theorem.

- Since $f \in C(0, \infty)$ and $\ell > 0$, it follows that the mapping $v \mapsto \rho(x)f(v + \ell)$ is continuous in $[0, \infty)$.
- From $\rho(x)\frac{f(v+\ell)}{v} = \rho(x)\frac{f(v+\ell)}{v+\ell}\frac{v+\ell}{v}$, using positivity of ρ and (f1) we deduce that the function $v \mapsto \rho(x)\frac{v+\ell}{v}$ is decreasing on $(0, \infty)$.
- For all $v \geq 0$, since $\rho \in L^\infty_{\text{loc}}(\mathbb{R}^N)$, we obtain that $\rho \in L^\infty(B(0, k))$, so the condition (5) is satisfied.
- By $\lim_{v \rightarrow \infty} f(v+\ell)/(v+1) = 0$ and $f \in C(0, \infty)$, there exists $M > 0$ such that $f(v+\ell) \leq M(v+1)$ for all $v \geq 0$. Therefore $\rho(x)f(v+\ell) \leq \|\rho\|_{L^\infty(B(0, k))}M(v+1)$ for all $v \geq 0$.
- We have

$$a_0(x) = \lim_{v \searrow 0} \frac{\rho(x)f(v+\ell)}{v} = +\infty$$

and

$$a_\infty(x) = \lim_{v \rightarrow \infty} \frac{\rho(x)f(v+\ell)}{v} = \lim_{v \rightarrow \infty} \rho(x) \frac{f(v+\ell)}{v+\ell} \cdot \frac{v+\ell}{v} = 0.$$

Thus, by Theorem 1 in [4], Problem (10) has a unique solution v_k which, by the maximum principle, is positive in $|x| < k$. Then $u_k = v_k + \ell$ satisfies (9). Define $u_k = \ell$ for $|x| > k$. The maximum principle implies that $\ell \leq u_k \leq u_{k+1}$ in $\mathbb{R}^N$.

We now justify the existence of a continuous function $v : \mathbb{R}^N \rightarrow \mathbb{R}$, $v > \ell$, such that $u_k \leq v$ in $\mathbb{R}^N$. We first construct a positive radially symmetric function w such that $-\Delta w = \Phi(r)$ ($r = |x|$) in $\mathbb{R}^N$ and $\lim_{r \rightarrow \infty} w(r) = 0$. A straightforward computation shows that

$$w(r) = K - \int_0^r \zeta^{1-N} \int_0^\zeta \sigma^{N-1} \Phi(\sigma) d\sigma d\zeta,$$

where

$$K = \int_0^\infty \zeta^{1-N} \int_0^\zeta \sigma^{N-1} \Phi(\sigma) d\sigma d\zeta,$$

provided the integral is finite. An integration by parts yields

$$\begin{aligned} \int_0^r \zeta^{1-N} \int_0^\zeta \sigma^{N-1} \Phi(\sigma) d\sigma d\zeta &= -\frac{1}{N-2} \int_0^r \frac{d}{d\zeta} \zeta^{2-N} \int_0^\zeta \sigma^{N-1} \Phi(\sigma) d\sigma d\zeta \\ &= \frac{1}{N-2} \left(-r^{2-N} \int_0^r \sigma^{N-1} \Phi(\sigma) d\sigma + \int_0^r \zeta \Phi(\zeta) d\zeta \right) \\ &< \frac{1}{N-2} \int_0^\infty \zeta \Phi(\zeta) d\zeta < +\infty. \end{aligned}$$

Moreover, w is decreasing and satisfies $0 < w(r) < K$ for all $r \geq 0$. Let $v > \ell$ be a function such that $w(r) = m^{-1} \int_0^{v(r)-\ell} \frac{t}{f(t+\ell)} dt$, where $m > 0$ is chosen such that $Km \leq \int_0^m \frac{t}{f(t+\ell)} dt$.

Next, by L'Hôpital's rule for the case $\frac{\infty}{\infty}$ (see [15, Theorem 3, p. 319]) we have

$$\lim_{x \rightarrow \infty} \frac{\int_0^x \frac{t}{f(t+\ell)} dt}{x} = \lim_{x \rightarrow \infty} \frac{x}{f(x+\ell)} = \lim_{x \rightarrow \infty} \frac{x+\ell}{f(x+\ell)} \cdot \frac{x}{x+\ell} = +\infty.$$

This means that there exists $x_1 > 0$ such that $\int_0^x \frac{t}{f(t)} dt \geq Kx$ for all $x \geq x_1$. It follows that for any $m \geq x_1$ we have $Km \leq \int_0^m \frac{t}{f(t)} dt$.

Since w is decreasing, we obtain that v is a decreasing function, too. Then

$$\int_0^{v(r)-\ell} \frac{t}{f(t+\ell)} dt \leq \int_0^{v(0)-\ell} \frac{t}{f(t+\ell)} dt = mw(0) = mK \leq \int_0^m \frac{t}{f(t+\ell)} dt.$$

It follows that $v(r) \leq m + \ell$ for all $r > 0$.

From $w(r) \rightarrow 0$ as $r \rightarrow \infty$ we deduce that $v(r) \rightarrow \ell$ as $r \rightarrow \infty$.

By the choice of v we have

$$\nabla w = \frac{1}{M} \frac{v-\ell}{f(v)} \nabla v \quad \text{and} \quad \Delta w = \frac{1}{m} \frac{v-\ell}{f(v)} \Delta v + \frac{1}{m} \left(\frac{v-\ell}{f(v)} \right)' |\nabla v|^2.$$

Since the mapping $u \mapsto f(u)/u$ is decreasing on $(0, \infty)$ we deduce that

$$\Delta v < \frac{m}{v-\ell} f(v) \Delta w = -\frac{M}{v-\ell} f(v) \Phi(r) \leq -f(v) \Phi(r). \quad (11)$$

By (9), (11) and our hypothesis (f1), we obtain that $u_k(x) \leq v(x)$ for each $|x| \leq k$ and so, for all $x \in \mathbb{R}^N$.

In conclusion,

$$u_1 \leq u_2 \leq \dots \leq u_k \leq u_{k+1} \leq \dots \leq v,$$

with $v(x) \rightarrow \ell$ as $|x| \rightarrow \infty$. Thus, there exists a function $u \leq v$ such that $u_k \rightarrow u$ pointwise in $\mathbb{R}^N$. In particular, this shows that $u > \ell$ in $\mathbb{R}^N$ and $u(x) \rightarrow \ell$ as $|x| \rightarrow \infty$.

A standard bootstrap argument (see, e.g., [10]) shows that u is a classical solution of the problem (2).

To conclude the proof, it remains to show that the solution found above is unique. Suppose that u and v are solutions of (2). It is enough to show that $u \leq v$ or, equivalently, $\ln u(x) \leq \ln v(x)$, for any $x \in \mathbb{R}^N$. Arguing by contradiction, there exists $\bar{x} \in \mathbb{R}$ such that $u(\bar{x}) > v(\bar{x})$. Since $\lim_{|x| \rightarrow \infty} (\ln u(x) - \ln v(x)) = 0$, we deduce that $\max_{\mathbb{R}^N} (\ln u(x) - \ln v(x))$ exists and is positive. At this point, say x_0 , we have

$$\nabla (\ln u(x_0) - \ln v(x_0)) = 0, \quad (12)$$

so

$$\frac{\nabla u(x_0)}{u(x_0)} = \frac{\nabla v(x_0)}{v(x_0)}. \quad (13)$$

By (f1) we obtain

$$\frac{f(u(x_0))}{u(x_0)} < \frac{f(v(x_0))}{v(x_0)}.$$

So, by (12) and (13),

$$\begin{aligned}
0 &\geq \Delta(\ln u(x_0) - \ln v(x_0)) \\
&= \frac{1}{u(x_0)} \cdot \Delta u(x_0) - \frac{1}{v(x_0)} \cdot \Delta v(x_0) - \frac{1}{u^2(x_0)} \cdot |\nabla u(x_0)|^2 + \frac{1}{v^2(x_0)} \cdot |\nabla v(x_0)|^2 \\
&= \frac{\Delta u(x_0)}{u(x_0)} - \frac{\Delta v(x_0)}{v(x_0)} = -\rho(x_0) \left(\frac{f(u(x_0))}{u(x_0)} - \frac{f(v(x_0))}{v(x_0)} \right) > 0,
\end{aligned}$$

which is a contradiction. Hence $u \leq v$ and the proof is concluded. $\square$

3 Proof of Theorem 2

3.1 Existence

Since f is an increasing positive function on $(0, \infty)$, there exists and is finite $\lim_{u \searrow 0} f(u)$, so f can be extended by continuity at the origin. Consider the Dirichlet problem

$$\begin{cases} -\Delta u_k = \rho(x)f(u_k) , & \text{if } |x| < k \\ u_k(x) = 0, & \text{if } |x| = k. \end{cases} \quad (14)$$

Using the same arguments as in case $\ell > 0$ we deduce that conditions (4) and (5) are satisfied. In what concerns assumption (6), we use both assumptions (f1) and (f2). Hence $f(u) \leq f(1)$ if $u \leq 1$ and $f(u)/u \leq f(1)$ if $u \geq 1$. Therefore $f(u) \leq f(1)(u+1)$ for all $u \geq 0$, which proves (6). The existence of a solution for (14) follows from (7) and (8). These conditions are direct consequences of our assumptions $\lim_{u \rightarrow \infty} f(u)/u = 0$ and $\lim_{u \searrow 0} f(u)/u = +\infty$. Thus, by the Brezis-Oswald theorem, Problem (14) has a unique solution. Define $u_k(x) = 0$ for $|x| > k$. Using the same arguments as in case $\ell > 0$, we obtain $u_k \leq u_{k+1}$ in $\mathbb{R}^N$.

Next, we prove the existence of a continuous function $v : \mathbb{R}^N \rightarrow \mathbb{R}$ such that $u_k \leq v$ in $\mathbb{R}^N$. We first construct a positive radially symmetric function w satisfying $-\Delta w = \Phi(r)$ ($r = |x|$) in $\mathbb{R}^N$ and $\lim_{r \rightarrow \infty} w(r) = 0$. We obtain

$$w(r) = K - \int_0^r \zeta^{1-N} \int_0^\zeta \sigma^{N-1} \Phi(\sigma) d\sigma d\zeta ,$$

where

$$K = \int_0^\infty \zeta^{1-N} \int_0^\zeta \sigma^{N-1} \Phi(\sigma) d\sigma d\zeta , \quad (15)$$

provided the integral is finite. By integration by parts we have

$$\begin{aligned}
\int_0^r \zeta^{1-N} \int_0^\zeta \sigma^{N-1} \Phi(\sigma) d\sigma d\zeta &= -\frac{1}{N-2} \int_0^r \frac{d}{d\zeta} \zeta^{2-N} \int_0^\zeta \sigma^{N-1} \Phi(\sigma) d\sigma d\zeta = \\
\frac{1}{N-2} \left(-r^{2-N} \int_0^r \sigma^{N-1} \Phi(\sigma) d\sigma + \int_0^r \zeta \Phi(\zeta) d\zeta \right) &< \frac{1}{N-2} \int_0^\infty \zeta \Phi(\zeta) d\zeta < \infty .
\end{aligned} \quad (16)$$

Therefore

$$w(r) < \frac{1}{N-2} \cdot \int_0^\infty \zeta \Phi(\zeta) d\zeta, \quad \text{for all } r > 0.$$

Let v be a positive function such that $w(r) = c^{-1} \int_0^{v(r)} t/f(t) dt$, where $c > 0$ is chosen such that $Kc \leq \int_0^c t/f(t) dt$. We argue in what follows that we can find $c > 0$ with this property. Indeed, by L'Hôpital's rule,

$$\lim_{x \rightarrow \infty} \frac{\int_0^x \frac{t}{f(t)} dt}{x} = \lim_{x \rightarrow \infty} \frac{x}{f(x)} = +\infty.$$

This means that there exists $x_1 > 0$ such that $\int_0^x t/f(t) dt \geq Kx$ for all $x \geq x_1$. It follows that for any $c \geq x_1$ we have $Kc \leq \int_0^c t/f(t) dt$.

On the other hand, since w is decreasing, we deduce that v is a decreasing function, too. Hence

$$\int_0^{v(r)} \frac{t}{f(t)} dt \leq \int_0^{v(0)} \frac{t}{f(t)} dt = c \cdot w(0) = c \cdot K \leq \int_0^c \frac{t}{f(t)} dt.$$

It follows that $v(r) \leq c$ for all $r > 0$.

From $w(r) \rightarrow 0$ as $r \rightarrow \infty$ we deduce that $v(r) \rightarrow 0$ as $r \rightarrow \infty$.

By the choice of v we have

$$\nabla w = \frac{1}{c} \cdot \frac{v}{f(v)} \nabla v \quad \text{and} \quad \Delta w = \frac{1}{c} \frac{v}{f(v)} \Delta v + \frac{1}{c} \left(\frac{v}{f(v)} \right)' |\nabla v|^2. \quad (17)$$

Combining the fact that $f(u)/u$ is a decreasing function on $(0, \infty)$ with relation (17), we deduce that

$$\Delta v < c \frac{f(v)}{v} \Delta w = -c \frac{f(v)}{v} \Phi(r) \leq -f(v) \Phi(r). \quad (18)$$

By (14) and (18) and using our hypothesis (f2), as already done for proving the uniqueness in the case $\ell > 0$, we obtain that $u_k(x) \leq v(x)$ for each $|x| \leq k$ and so, for all $x \in \mathbb{R}^N$.

We have obtained a bounded increasing sequence

$$u_1 \leq u_2 \leq \dots \leq u_k \leq u_{k+1} \leq \dots \leq v,$$

with v vanishing at infinity. Thus, there exists a function $u \leq v$ such that $u_k \rightarrow u$ pointwise in $\mathbb{R}^N$. A standard bootstrap argument implies that u is a classical solution of the problem (2).

3.2 Uniqueness

We split the proof into two steps. Assume that u_1 and u_2 are solutions of Problem (2). We first prove that if $u_1 \leq u_2$ then $u_1 = u_2$ in $\mathbb{R}^N$. In the second step we find a positive solution $u \leq \min\{u_1, u_2\}$ and thus, using the first step, we deduce that $u = u_1$ and $u = u_2$, which proves the uniqueness.

STEP I. We show that $u_1 \leq u_2$ in $\mathbb{R}^N$ implies $u_1 = u_2$ in $\mathbb{R}^N$. Indeed, since

$$u_1 \Delta u_2 - u_2 \Delta u_1 = \rho(x) u_1 u_2 \left(\frac{f(u_2)}{u_2} - \frac{f(u_1)}{u_1} \right) \geq 0,$$

it is sufficient to check that

$$\int_{\mathbb{R}^N} (u_1 \Delta u_2 - u_2 \Delta u_1) = 0 \quad (19)$$

Let $\psi \in C_0^\infty(\mathbb{R}^N)$ be such that $\psi(x) = 1$ for $|x| \leq 1$ and $\psi(x) = 0$ for $|x| \geq 2$, and denote $\psi_n := \psi(x/n)$ for any positive integer n . Set

$$I_n := \int_{\mathbb{R}^N} (u_1 \Delta u_2 - u_2 \Delta u_1) \psi_n dx.$$

We claim that $I_n \rightarrow 0$ as $n \rightarrow \infty$. Indeed,

$$|I_n| \leq \int_{\mathbb{R}^N} |u_1 \Delta u_2| \psi_n dx + \int_{\mathbb{R}^N} |u_2 \Delta u_1| \psi_n dx.$$

So, by symmetry, it is enough to prove that $J_n := \int_{\mathbb{R}^N} |u_1 \Delta u_2| \psi_n dx \rightarrow 0$ as $n \rightarrow \infty$. But, from (2),

$$\begin{aligned} J_n &= \int_{\mathbb{R}^N} |u_1 f(u_2) \rho(x)| \psi_n dx = \int_n^{2n} \int_{|x|=r} |u_1(x) f(u_2(x)) \rho(x)| dx dr \\ &\leq \int_n^{2n} \Phi(r) \int_{|x|=r} |u_1(x) f(u_2(x))| dx dr \leq \int_n^{2n} \Phi(r) \int_{|x|=r} |u_1(x)| M(u_2 + 1) dx dr. \end{aligned} \quad (20)$$

Since $u_1(x), u_2(x) \rightarrow 0$ as $|x| \rightarrow \infty$, we deduce that u_1 and u_2 are bounded in $\mathbb{R}^N$. Returning to (20) we have

$$J_n \leq M(\|u_2\|_{L^\infty(\mathbb{R}^N)} + 1) \sup_{|x| \geq n} |u_1(x)| \cdot \frac{\omega_N}{N} \int_n^{2n} \Phi(r) r^{N-1} dr \leq C \int_0^\infty \Phi(r) r^{N-1} dr \cdot \sup_{|x| \geq n} |u_1(x)|.$$

Since $u_1(x) \rightarrow 0$ as $|x| \rightarrow \infty$, we have $\sup_{|x| \geq n} |u_1(x)| \rightarrow 0$ as $n \rightarrow \infty$ which shows that $J_n \rightarrow 0$. In particular, this implies $I_n \rightarrow 0$ as $n \rightarrow \infty$.

We recall in what follows the Lebesgue Dominated Convergence Theorem (see [2, Theorem IV.2]).

Theorem 3. *Let $f_n : \mathbb{R}^N \rightarrow \mathbb{R}$ be a sequence of functions in $L^1(\mathbb{R}^N)$. We assume that*

(i) $f_n(x) \rightarrow f(x)$ a.e. in $\mathbb{R}^N$,

(ii) *there exists $g \in L^1(\mathbb{R}^N)$ such that, for all $n \geq 1$, $|f_n(x)| \leq g(x)$ a.e. in $\mathbb{R}^N$.*

Then $f \in L^1(\mathbb{R}^N)$ and $\|f_n - f\|_{L^1} \rightarrow 0$ as $n \rightarrow \infty$.

Taking $f_n := (u_1 \Delta u_2 - u_2 \Delta u_1) \psi_n$ we deduce $f_n(x) \rightarrow u_1(x) \Delta u_2(x) - u_2(x) \Delta u_1(x)$ as $n \rightarrow \infty$. To apply Theorem 3 we need to show that $u_1 \Delta u_2 - u_2 \Delta u_1 \in L^1(\mathbb{R}^N)$. For this purpose it is sufficient to prove that $u_1 \Delta u_2 \in L^1(\mathbb{R}^N)$. Indeed,

$$\int_{\mathbb{R}^N} |u_1 \Delta u_2| \leq \|u_1\|_{L^\infty(\mathbb{R}^N)} \int_{\mathbb{R}^N} |\Delta u_2| = C \int_{\mathbb{R}^N} |\rho(x) f(u_2)|.$$

Thus, using $f(u) \leq f(1)(u + 1)$ and since u_2 is bounded, the above inequality yields

$$\begin{aligned} \int_{\mathbb{R}^N} |u_1 \Delta u_2| &\leq C \int_{\mathbb{R}^N} |\rho(x)(u_2 + 1)| \\ &\leq C \int_0^\infty \int_{|x|=r} \Phi(r) dx dr \leq C \int_0^\infty \Phi(r) r^{N-1} < +\infty. \end{aligned}$$

This shows that $u_1 \Delta u_2 \in L^1(\mathbb{R}^N)$ and the proof of Step I is completed.

STEP II. Let u_1, u_2 be arbitrary solutions of Problem (2). For all integer $k \geq 1$, denote $\Omega_k := \{x \in \mathbb{R}^N; |x| < k\}$. The Brezis-Oswald theorem implies that the problem

$$\begin{cases} -\Delta v_k = \rho(x)f(v_k) & \text{in } \Omega_k \\ v_k = 0 & \text{on } \partial\Omega_k \end{cases}$$

has a unique solution $v_k \geq 0$. Moreover, by the Maximum Principle, $v_k > 0$ in Ω_k . We define $v_k = 0$ for $|x| > k$. Applying again the Maximum Principle we deduce that $v_k \leq v_{k+1}$ in $\mathbb{R}^N$. Now we prove that $v_k \leq u_1$ in $\mathbb{R}^N$, for all $k \geq 1$. Obviously, this happens outside Ω_k . On the other hand

$$\begin{cases} -\Delta u_1 = \rho(x)f(u_1) & \text{in } \Omega_k \\ u_1 > 0 & \text{on } \partial\Omega_k \end{cases}$$

Arguing by contradiction, we assume that there exists $\bar{x} \in \Omega_k$ such that $v_k(\bar{x}) > u_1(\bar{x})$. Consider the function $h : \Omega_k \rightarrow \mathbb{R}$, $h(x) = \ln v_k(x) - \ln u_1(x)$. Since u_1 is bounded in Ω_k and $\inf_{\partial\Omega_k} u_1 > 0$ we have $\lim_{|x| \rightarrow k} h(x) = -\infty$. We deduce that $\max_{\Omega_k} (\ln v_k(x) - \ln u_1(x))$ exists and is positive. Using the same

argument as in the case $\ell > 0$ we deduce that $v_k \leq u_1$ in Ω_k , so in $\mathbb{R}^N$. Similarly we obtain $v_k \leq u_2$ in $\mathbb{R}^N$. Hence $v_k \leq \bar{u} := \min\{u_1, u_2\}$. Therefore $v_k \leq v_{k+1} \leq \dots \leq \bar{u}$. Thus there exists a function u such that $v_k \rightarrow u$ pointwise in $\mathbb{R}^N$. Repeating a previous argument we deduce that $u \leq \bar{u}$ is a classical solution of Problem (2). Moreover, since $u \geq v_k > 0$ in Ω_k and for all $k \geq 1$, we deduce that $u > 0$ in $\mathbb{R}^N$. This concludes the proof of Step II.

Combining Steps I and II we conclude that $u_1 = u_2$ in $\mathbb{R}^N$. □

References

- [1] M. Aizenman and B. Simon, Brownian motion and Harnack inequality for Schrödinger operators, *Comm. Pure Appl. Math.* **35** (1982), 209-273.
- [2] H. Brezis, *Analyse Fonctionnelle. Théorie et applications*, Masson, Paris, 1983.
- [3] H. Brezis and S. Kamin, Sublinear elliptic equations in $\mathbb{R}^n$, *Manuscripta Math.* **74** (1992), 87-106.
- [4] H. Brezis and L. Oswald, Remarks on sublinear elliptic equations, *Nonlinear Anal., T.M.A.* **10** (1986), 55-64.
- [5] T.-L. Dinu, Entire solutions of the nonlinear eigenvalue logistic problem with sign-changing potential and absorption, *Commun. Pure Appl. Anal.* **2** (2003), 311-321.
- [6] A. Edelson, Asymptotic properties of semilinear equations, *Canad. Math. Bull.* **32** (1989), 34-46.
- [7] A. Edelson, Entire solutions of singular elliptic equations, *J. Math. Anal. Appl.* **139** (1989), 523-532.
- [8] H. Egnell, Asymptotic results for finite energy solutions of semilinear elliptic equations, *J. Differential Equations* **98** (1992), 34-56.

- [9] N. Fukagai, Existence and uniqueness of entire solutions of second order sublinear elliptic equations, *Funkcial. Ekvac.* **29** (1986), 151-165.
- [10] D. Gilbarg and N. Trudinger, *Elliptic partial differential equations of second order*, Classics in Mathematics, Springer-Verlag, Berlin, 2001.
- [11] N. Kawano, On bounded entire solutions of semilinear elliptic equations, *Hiroshima Math. J.* **14** (1984), 125-158.
- [12] M. A. Krasnoselskii, *Positive Solutions of Operator Equations*, Noordhoff, Groningen, 1964.
- [13] K. El Mabrouk, Entire bounded solutions for a class of sublinear elliptic equations, *Nonlinear Anal.* **58** (2004), 205-218.
- [14] M. Naito, A note on bounded positive entire solutions of semilinear elliptic equations, *Hiroshima Math. J.* **14** (1984), 211-214.
- [15] M. Nicolescu, N. Dinculeanu and S. Marcus, *Mathematical Analysis I*, Ed. Didact. Pedag., Bucureşti, 1971 [in Romanian].
- [16] B. O'Neill, *Semi-Riemannian geometry with applications to relativity*, Pure and Applied Mathematics, vol. 103, Academic Press, New York, 1983.